

Two-Loop $\mathcal{O}(\alpha_s^2)$ Correction to the $H \rightarrow b\bar{b}$ Decay Rate Induced by the Top Quark

Bernd A. Kniehl

Max-Planck-Institut für Physik, Werner-Heisenberg-Institut
Föhringer Ring 6, 80805 Munich, Germany

Abstract

We present the two-loop QED and QCD corrections to the $f\bar{f}H$ Yukawa coupling that are induced by the exchange of a virtual photon or gluon, respectively, with a heavy-fermion loop inserted. As an application, we study the corresponding $\mathcal{O}(\alpha_s^2 M_H^2/m_t^2)$ correction to the $H \rightarrow b\bar{b}$ decay rate.

A Higgs boson with $M_H \lesssim 135$ GeV decays dominantly to $b\bar{b}$ pairs [1]. This decay mode will be of prime importance for Higgs-boson searches at LEP 2 [2], the Tevatron [3]—or a possible 4-TeV upgrade thereof [4]—, and the next e^+e^- linear collider [5]. Techniques for the measurement of the $H \rightarrow b\bar{b}$ branching fraction at a $\sqrt{s} = 500$ GeV e^+e^- linear collider have been elaborated in Ref. [6].

The present knowledge of quantum corrections to the $H \rightarrow b\bar{b}$ decay rate has been reviewed very recently in Ref. [1]. The full one-loop electroweak corrections to this process are well established [7, 8]. At two loops, the universal [9] and non-universal [10] $\mathcal{O}(\alpha_s G_F m_t^2)$ corrections have recently been calculated. The pure QCD corrections are most significant numerically. In $\mathcal{O}(\alpha_s)$, their full m_b dependence is known [11]. In $\mathcal{O}(\alpha_s^2)$, the first [12] and second [13] terms of the expansion in m_b^2/M_H^2 have been found.

The results of Refs. [12, 13] take into account five quark flavours. In $\mathcal{O}(\alpha_s^2)$, there are additional contributions involving a virtual top quark, which are not formally suppressed. These corrections may be viewed as the absorptive parts of the three-loop Higgs-boson self-energy diagrams that involve a top-quark loop, a bottom-quark loop, and two gluon lines. These diagrams may be divided in two classes: the so-called double-triangle diagrams and the diagrams where a gluon line with a top-quark loop insertion is attached in all possible ways to the one-loop seed diagram involving the bottom quark. The first class has been considered just recently [14]. In this letter, we shall study the second class.

We shall proceed along the lines of Ref. [15],¹ where the two-loop $f\bar{f}\gamma$ vertex correction due to a virtual massive fermion, F , was derived in QED, assuming $m_F^2 \ll |s|$, where s

¹We take this opportunity to correct two misprints in the published version of Ref. [15], which are absent in the preprint. In the second line of Eq. (5), the terms $-\varphi^2$ and $53/2$ should be replaced by $+\varphi^2$ and $53/3$.

is the photon invariant mass squared. In this limit, only the Dirac Form factor, which multiplies γ^μ , survives. It has the form [15]

$$\Gamma^\mu = \gamma^\mu \left[1 + \frac{\alpha}{\pi} Q_f^2 F_1(s) + \left(\frac{\alpha}{\pi} \right)^2 Q_f^2 N_F Q_F^2 F_2^{(F)}(s) + \dots \right], \quad (1)$$

where Q_f is the electric charge of f (in units of the positron charge), $N_f = 1$ (3) for leptons (quarks), and the dots represent other contributions in $\mathcal{O}(\alpha^2)$ and higher orders. By adjusting coupling constants and colour factors, one immediately obtains the corresponding QCD expansion for the case when f and F are quarks. Specifically, one substitutes $\alpha \rightarrow \alpha_s$, $Q_f^2 \rightarrow C_F = (N_c^2 - 1)/(2N_c) = 4/3$, and $Q_F^2 \rightarrow T = 1/2$, where $N_c = 3$ is number of colours and $\text{tr } t^a t^b = T \delta^{ab}$, with t^a ($a = 1, \dots, 8$) being the generators of the quark (defining) representation of $\text{SU}(N_c)$. In particular, the m_t -dependent $\mathcal{O}(\alpha_s^2)$ correction to the non-singlet contribution to $R = \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$ is given by

$$\delta R_{NS} = 2 \left(\frac{\alpha_s}{\pi} \right)^2 C_F T \text{Re } F_2^{(F)}(s), \quad (2)$$

which is in agreement with Ref. [16].

Similarly to Eq. (1), the QED expansion of the $f\bar{f}H$ coupling may be written as

$$\Gamma = 1 + \frac{\alpha}{\pi} Q_f^2 H_1(s) + \left(\frac{\alpha}{\pi} \right)^2 Q_f^2 N_F Q_F^2 H_2^{(F)}(s) + \dots \quad (3)$$

We have $H_1(s) = (1/4)\hat{\delta}_{em}|_{h=s}$, where $\hat{\delta}_{em}$ may be found in Eq. (2.18) of Ref. [8]. $H_1(s)$ is plagued by an infrared singularity, which is cancelled by a similar contribution from soft-photon bremsstrahlung when a physical observable, such as the $H \rightarrow f\bar{f}$ decay rate, is computed. In Ref. [8], the infrared singularity is regularized by an infinitesimal photon mass, μ . Up to terms proportional to m_f^2/s , one has

$$H_1(s) = \frac{L}{2}(z-1) - \frac{z^2}{4} + \frac{7}{2}\zeta(2) - \frac{1}{2}, \quad (4)$$

where $\zeta(2) = \pi^2/6$, $L = \ln(\mu^2/m_f^2)$, and $z = \ln(-s/m_f^2 - i\epsilon)$. In the following, we shall need an expression for $H_1(s)$ appropriate for a neutral gauge boson with arbitrary mass, $\sqrt{\sigma}$. From Eq. (2.6) of Ref. [8] one extracts

$$H_1(s, \sigma) = \frac{1}{2} \left(\text{Li}_2 \left(1 + \frac{s+i\epsilon}{\sigma} \right) - \zeta(2) \right), \quad (5)$$

where Li_2 denotes the dilogarithm.

The Feynman diagrams contributing to $H_2^{(F)}(s)$ are depicted in Fig. 1. $H_2^{(F)}(s)$ is infrared-finite and devoid of mass singularities associated with f . It may be conveniently calculated by convoluting $H_1(s, \sigma)$ with the imaginary part of the one-loop contribution of F to the photon self-energy,

$$\text{Im } \Pi_{AA}^{(F)}(s) = \frac{\alpha}{3} N_F Q_F^2 s P(s), \quad (6)$$

where

$$P(s) = \left(1 + \frac{2m_F^2}{s}\right) \sqrt{1 - \frac{4m_F^2}{s}}. \quad (7)$$

The precise relation reads

$$H_2^{(F)}(s) = \frac{1}{3} \int_{4m_F^2}^{\infty} \frac{d\sigma}{\sigma} P(\sigma) H_1(s, \sigma). \quad (8)$$

After a straightforward calculation, one finds

$$\begin{aligned} 3H_2^{(F)}(s) &= \text{Li}_3(-\rho_-^2) + \frac{1}{3} \left(5 - \frac{1}{r}\right) \sqrt{1 + \frac{1}{r}} \left(\text{Li}_2(-\rho_-^2) + \varphi^2 + \frac{\zeta(2)}{2}\right) - \frac{2}{3} \varphi^3 - \zeta(2) \varphi \\ &\quad + \frac{2}{3} \left(-\frac{14}{3} + \frac{1}{r}\right) \gamma - \zeta(3) + \frac{82}{27} - \frac{2}{3r}, \quad r \leq -1, \\ &= \text{Cl}_3(2\Phi) - \frac{1}{3} \left(5 - \frac{1}{r}\right) \sqrt{-\frac{1}{r} - 1} \text{Cl}_2(2\Phi) \\ &\quad + \frac{2}{3} \left(-\frac{14}{3} + \frac{1}{r}\right) \gamma - \zeta(3) + \frac{82}{27} - \frac{2}{3r}, \quad -1 \leq r \leq 0, \\ &= \text{Li}_3(r_-^2) + \frac{1}{3} \left(5 - \frac{1}{r}\right) \sqrt{1 + \frac{1}{r}} \left(\text{Li}_2(r_-^2) + f^2 - \zeta(2)\right) - \frac{2}{3} f^3 + 2\zeta(2)f \\ &\quad + \frac{2}{3} \left(-\frac{14}{3} + \frac{1}{r}\right) g - \zeta(3) + \frac{82}{27} - \frac{2}{3r} \\ &\quad + i\pi \left[f^2 - \frac{1}{3} \left(5 - \frac{1}{r}\right) \sqrt{1 + \frac{1}{r}} f + \frac{14}{9} - \frac{1}{3r} \right], \quad r \geq 0, \end{aligned} \quad (9)$$

where $\zeta(3) = 1.20205690315959428540\dots$, Li_3 is the trilogarithm, Cl_2 (Cl_3) is the (generalized) Clausen function of second (third) order,

$$\begin{aligned} r &= \frac{s}{4m_F^2}, \quad \rho_{\pm} = \sqrt{-r} \pm \sqrt{-r-1}, \quad r_{\pm} = \sqrt{1+r} \pm \sqrt{r}, \\ \varphi &= \ln \rho_+ = \text{arcosh} \sqrt{-r}, \quad \Phi = \arcsin \sqrt{-r}, \quad f = \ln r_+ = \text{arsinh} \sqrt{r}, \\ \gamma &= \ln(\rho_+ + \rho_-) = \ln(2\sqrt{-r}), \quad g = \ln(r_+ - r_-) = \ln(2\sqrt{r}). \end{aligned} \quad (10)$$

Note that $H_1(s)$ and $H_2^{(F)}(s)$ develop imaginary parts above the $f\bar{f}$ -pair production threshold, i.e., for $s > 4m_f^2 = 0$.

It is instructive to study the limiting behaviour of $H_2^{(F)}(s)$. For $s \rightarrow -\infty$ and $s \rightarrow -0$, one has

$$\begin{aligned} 3H_2^{(F)}(s) &= -\frac{2}{3} \gamma^3 + \frac{5}{3} \gamma^2 - \left(\zeta(2) + \frac{28}{9}\right) \gamma - \zeta(3) + \frac{5}{6} \zeta(2) + \frac{82}{27} + \frac{3\gamma}{2r} \\ &\quad + \mathcal{O}\left(\frac{\gamma^2}{r^2}\right), \quad r \ll -1, \\ &= \frac{r}{5} \left(-4\gamma + \frac{107}{15}\right) + \frac{r^2}{35} \left(6\gamma - \frac{529}{70}\right) + \mathcal{O}(r^3\gamma), \quad -1 \ll r \leq 0, \end{aligned} \quad (11)$$

respectively. The corresponding expansions for positive s may be found by analytic continuation, i.e., by substituting $\gamma = g - i\pi/2$. In compliance with the Appelquist-Carazzone theorem [17], the loop fermion, F , decouples for $m_F^2 \gg |s|$.

In Fig. 2, $\text{Re } H_2^{(F)}(s)$ is plotted as a function of $r = s/(4m_F^2)$. At $r \approx 5.62$, $\text{Re } H_2^{(F)}(s)$ assumes its maximum value, 0.432. Its expansions, which emerge from Eq. (11) through analytic continuation, are also shown. Obviously, they provide an excellent approximation for $r \gtrsim 1$ and $r \lesssim 1$, respectively.

The QCD expansion of the $f\bar{f}H$ coupling for the case when f and F are quarks may be obtained from Eq. (3) through the substitutions specified below Eq. (1). As an application, we consider the m_t -dependent $\mathcal{O}(\alpha_s^2)$ correction to the $H \rightarrow b\bar{b}$ decay rate arising from the Feynman diagrams in Fig. 1 with $f = b$, $F = t$, and the photon replaced by a gluon. Similarly to Eq. (2), the relative shift is

$$\begin{aligned} \frac{\delta\Gamma(H \rightarrow b\bar{b})}{\Gamma(H \rightarrow b\bar{b})} &= 2 \left(\frac{\alpha_s}{\pi} \right)^2 C_F T \text{Re } H_2^{(t)}(M_H^2) \\ &= \left(\frac{\alpha_s}{\pi} \right)^2 \left[\frac{M_H^2}{45m_t^2} \left(2 \ln \frac{m_t^2}{M_H^2} + \frac{107}{15} \right) + \mathcal{O} \left(\frac{M_H^4}{m_t^4} \ln \frac{m_t^2}{M_H^2} \right) \right], \end{aligned} \quad (12)$$

where the second line is appropriate for $M_H \lesssim 2M_W$, where the $H \rightarrow b\bar{b}$ decay rate is most relevant phenomenologically. The coefficient of $(\alpha_s/\pi)^2$ in Eq. (12) ranges between 2.38×10^{-2} for $(M_H, m_t) = (60, 200)$ GeV and 0.502 for $(M_H, m_t) = (1000, 150)$ GeV; its value at $M_H = 2m_t$ is 0.353. This has to be compared with the value 29.14671 due to five massless quark flavours [12]. On the other hand, the finite- m_b term in $\mathcal{O}(\alpha_s^2)$ has the coefficient $-87.72459 (\overline{m}_b(M_H)/M_H)^2$ [13], where $\overline{m}_b(\mu)$ is the bottom-quark $\overline{\text{MS}}$ mass at renormalization scale μ . Assuming $\alpha_s(M_Z) = 0.118$ [18] and $m_b = 4.72$ GeV [19], this amounts to -0.200 (-2.37×10^{-2}) at $M_H = 60$ GeV ($2M_W$). In conclusion, the $\mathcal{O}(\alpha_s^2 M_H^2/m_t^2)$ correction to $\Gamma(H \rightarrow b\bar{b})$ arising from the Feynman diagrams shown in Fig. 1 is comparable in size with the $\mathcal{O}(\alpha_s^2 m_b^2/M_H^2)$ correction, but has the opposite sign.

References

- [1] B.A. Kniehl, Phys. Rep. **240** (1994) 211.
- [2] E. Gross, B.A. Kniehl, and G. Wolf, Z. Phys. **C 63** (1994) 417.
- [3] A. Stange, W. Marciano, and S. Willenbrock, Phys. Rev. **D 49** (1994) 1354.
- [4] J.F. Gunion and T. Han, UCD Report No. UCD-94-10, hep-ph/9404244 (April 1994); A. Stange, W. Marciano, and S. Willenbrock, Report No. BNL-60340 and ILL-(TH)-94-8, hep-ph/9404247 (April 1994).
- [5] V. Barger, K. Cheung, B.A. Kniehl, and R.J.N. Phillips, Phys. Rev. **D 46** (1992) 3725; V. Barger, K. Cheung, A. Djouadi, B.A. Kniehl, and P.M. Zerwas, Phys. Rev.

- D 49** (1994) 79; D. Haidt et al., in *Proceedings of the Workshop on e^+e^- Collisions at 500 GeV: The Physics Potential*, Munich, Annecy, Hamburg, ed. by P.M. Zerwas, Part A, DESY Report DESY 92-123A (August 1992) p. 1 and Part C, DESY Report DESY 93-123C (December 1993) p. 1; E. Boos, M. Sachwitz, H.J. Schreiber, and S. Shichanin, Z. Phys. **C 61** (1994) 675.
- [6] M.D. Hildreth, T.L. Barklow, and D.L. Burke, Phys. Rev. **D 49** (1994) 3441.
- [7] J. Fleischer and F. Jegerlehner, Phys. Rev. **D 23** (1981) 2001; D.Yu. Bardin, B.M. Vilenskiĭ, P.Kh. Khristov, Yad. Fiz. **53** (1991) 240 [Sov. J. Nucl. Phys. **53** (1991) 152]; A. Dabelstein and W. Hollik, Z. Phys. **C 53** (1992) 507.
- [8] B.A. Kniehl, Nucl. Phys. **B 376** (1992) 3.
- [9] B.A. Kniehl, Phys. Rev. **D 50** (1994) 3314.
- [10] A. Kwiatkowski and M. Steinhauser, Phys. Lett. **B 338** (1994) 66; B.A. Kniehl and M. Spira, DESY Report No. 94-102 (October 1994).
- [11] E. Braaten and J.P. Leveille, Phys. Rev. **D 22** (1980) 715; N. Sakai, Phys. Rev. **D 22** (1980) 2220; T. Inami and T. Kubota, Nucl. Phys. **B 179** (1981) 171; M. Drees and K. Hikasa, Phys. Lett. **B 240** (1990) 455; (E) **B 262** (1990) 497.
- [12] S.G. Gorishny, A.L. Kataev, S.A. Larin, and L.R. Surguladze, Mod. Phys. Lett. **A 5** (1990) 2703; Phys. Rev. **D 43** (1991) 1633; A.L. Kataev and V.T. Kim, Mod. Phys. Lett. **A 9** (1994) 1309.
- [13] L.R. Surguladze, University of Oregon Report No. OITS 543, hep-ph/9405325 (May 1994).
- [14] K.G. Chetyrkin, J.H. Kühn, and A. Kwiatkowski, Karlsruhe University Report No. TTP94-10, hep-ph/9407271 (July 1994), to appear in *Proceedings of the 1994 Zeuthen Workshop on Elementary Particle Theory: Physics at LEP200 and Beyond*, Teupitz, April 10–15, 1994, ed. by J. Blümlein and T. Riemann, Nucl. Phys. B (Proc. Suppl.).
- [15] B.A. Kniehl, UW-Madison Report No. MAD/PH/545 (December 1989); Phys. Lett. **B 237** (1990) 127.
- [16] K.G. Chetyrkin, Phys. Lett. **B 307** (1993) 169; D.E. Soper and L.R. Surguladze, University of Oregon Report No. OITS 545, hep-ph/9406407 (June 1994).
- [17] T. Appelquist and J. Carazzone, Phys. Rev. **D 11** (1975) 2856.
- [18] S. Bethke, J. Phys. **G 17** (1991) 1455; to appear in *Proceedings of the Tennessee International Symposium on Radiative Corrections: Status and Outlook*, Gatlinburg, Tennessee, June 27–July 1, 1994, ed. by B.F.L. Ward (World Scientific, Singapore, 1994).
- [19] C.A. Dominguez and N. Paver, Phys. Lett. **B 293** (1992) 197.

FIGURE CAPTIONS

Figure 1: Feynman diagrams pertinent to the two-loop $f\bar{f}H$ vertex correction induced by a virtual heavy fermion, F .

Figure 2: $\text{Re } H_2^{(F)}(s)$ as a function of $r = s/(4m_F^2)$ [see Eq. (9)] and its expansions [see Eq. (11)].

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9410356v1>

This figure "fig2-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-ph/9410356v1>